AVROM E. HOWARD, MSc, PGeo, FGA

Professional Geologist & Certified Gemologist

GEOLOGICAL REPORT ON THE POWDERHORN LAKE PROPERTY

Springdale Area, Newfoundland, Canada

Prepared For: Champion Minerals Inc.



Prepared By: Avrom E. Howard, MSc, FGA, PGeo (Ontario)

March 1, 2007

TABLE OF CONTENTS

2.	Table of Contents	2			
3.	Summary				
4.	Introduction and Terms of Reference				
5.	Disclaimer				
6.	Property Description and Location 6.1 Location 6.2 Mineral Exploration Licences 6.3 Environmental Liabilities and Permits				
7.	Accessibility, Climate, Local Resources, Infrastructure and Physiography				
8.	History	14			
9.	Geological Setting 9.1 Regional Geology 9.2 Local Geology 9.3 Property Geology	18 18 20 21			
10.	Deposit Types	25			
11.	Mineralization 11.1 The Main Showing 11.2 The Road Showing 11.3 South Zone 11.4 Mineral Point (or Old Powderhorn) Zone 11.5 Dead Tree Zone 11.6 Pearl Zone 11.7 Unnamed Gossan	26 26 27 28 28 29 29			
12.	Exploration 12.1 2001 Field Season 12.2 2005-06 Field Season 12.3 Personnel	30 30 30 31			
13.	Drilling 13.1 2001 Drilling Program 13.2 2006 Drilling Program	32 32 35			
14.	14. Sampling Method and Approach				
15.	5. Sample Preparation, Analyses and Security				
16.	6. Data Verification				

17.	Adjacent Properties							
18.	Mineral Processing and Metallurgical Testing							
19.	Mineral I	Resource and Mineral Reserve Estimates	50					
20.	Other Relevant Data and Information							
21.	Interpretation and Conclusions							
22.	22.1	mendations Recommendations Budget						
23.	Reference	ees	56					
24.	Certificat	te Of The Author	61					
25.	5. Consent Of The Author							
26.	26. Certificate Of Champion's Qualified Person							
FIGI	URES AND	TABLES						
Figu	ıre 1:	Location of The Powderhorn Lake Property	8					
Figu	ıre 2:	License Location Map, Powderhorn Property	9					
Figure 3:		Compilation Map Showing Results Of An Induced Polarization Survey Conducted In 2000 and The Location Of Follow Up Drill Holes Completed In 2001 (From Dwyer, 2001 And Reed, 2000)	17					
Figure 4:		Tectonic Subdivisions Of Newfoundland (After Williams Et Al, 1988)	18					
Figure 5:		5: Map Showing The Location Of All Significant VHMS Deposits Found Within The Dunnage Zone (After Swinden, 1991)						
Figu	ıre 6:	Geology Of The Powderhorn Lake Area (From Dickson, 2000)	23					
Figure 7:		e 7: Geological Map Of The Powderhorn Lake Property (From Dwyer, 20						
Figure 8:		e 8: Cross-Section Of Drill Hole PH01-09 Showing Copper And Zinc (In Percent) And Gold (50 Ppb Or Greater) (From Dwyer, 2001)						
Figure 9:		Plan Map Showing The Location Of The 2006 Champion Drilling,	36					

Figure 10:	Cross Section A-A' (Facing Northwest)							
Figure 11:	Cross Section B-B' (Facing Northeast)							
Table One:	Summary Of The Powderhorn Lake Property Licences							
Table Two:	Selected Assay Data For Hole PH01-09							
Table Three:	2006	Drilling Program, Hole Location Coordinates	35					
Table Four: High		ghts Of Assay Results From The 2006 Drill Core Sampling	36					
Table Five:		ion of Duplicate Sample Analyses For Primary Metals erest (Error %)	47					
LIST OF APPENDICES								
Appendix One	e :	Drill Logs, 2006 Drilling	65					
Appendix Two):	Assay Certificates, 2006 Drilling	91					
Appendix Thre	ee:	Graphs Of Results For The Blank Standards	125					
Appendix Fou	ır:	Graphs Of Results For The Certified Reference Standard	130					
Appendix Five	:	Bias Plots For Each Of The Duplicate Sample Types For Each Of The Principal Metals	134					
Appendix Six:		Option Agreement Between Copper Hill Resources And Champion Natural Health.Com	148					
Appendix Sev	en:	Name Change from Champion Natural Health.Com, To Champion Minerals Inc.	211					

3. Summary

The Powderhom Lake Property, located in central Newfoundland, is held by Copper Hill Resources Inc. a company based in St. John's, Newfoundland. The property hosts two varieties of metallic mineral occurrences, in turn indicative of two distinct geologic settings: (1) volcanic-hosted massive sulphide ("VHMS") polymetallic mineralization, and (2) nickel sulphide mineralization associated with mafic intrusives. The VHMS style mineralization is hosted by the Roberts Arm Group, a regional geological unit that hosts a number of past-productive deposits, such as Buchans and Gullbridge. Selected hand samples taken by previous workers from the sulphide occurrences on the property have returned values of up to 4.5 percent zinc, 2.14 percent copper, 1.87 percent lead, and 1.2 grams per tonne ("g/t") gold. Diamond drilling in the Pearl Zone has intersected mineralization within a stratigraphic interval, with core samples returning values of up to 4 percent zinc over 0.6 meters ("m") and 3.1 percent zinc over 5 m (apparent thickness). More recent exploration efforts have focused on lithological, alteration, mineral zonation and geophysical features that may enable a trend or "vector" to be discerned, pointing toward the assumed VHMS source of this mineralization.

Occurrences of nickel sulphide mineralization on the property are restricted a gabbroic to dioritic intrusive suite. They do not appear to be of any economic significance.

In order to define possible VHMS targets on the property it is recommended that a primary geologic approach is taken in order to establish an observation based geologic model after which geophysical data may be incorporated in order to test and refine it, leading to the delineation of discrete drill targets. This approach should incorporate a fresh geological mapping campaign, with particular emphasis on stratigraphy, structure, alteration and mineral zonation. This should be followed by the re-logging of all drill core, incorporating the sub-surface geology into the established geological classifications and with a focus on any features that may indicate a direction toward better thicknesses and grades of sulphide mineralization. The combination of surface and subsurface geologic data should allow for the construction of a three-dimensional geologic model, upon which the existing and perhaps additional geophysical data may be incorporated to verify and perhaps refine the geologic model. Drill targets may then be selected, accordingly. A budget of C \$300,000 is estimated.